



Post-operative Physiotherapy Protocol Hip Arthroscopy

Ms Nichole Hamilton

**June 2007
(Revised 2011)**

Hip arthroscopy offers a method of treatment for hip disorders including (most commonly) labral tears, loose bodies, impinging osteophytes and chondral injuries. The benefit of hip arthroscopy is that the surgical technique is more accurate and less invasive, and the rehabilitation is much simpler than for open surgery.

The mechanism of injury for labral tear can be significant trauma involving hip rotation. Patients can also present with a gradual progressive onset of symptoms (without trauma) if a predisposition to increased load on the anterior hip and labrum is apparent. Predisposing factors that can increase stresses on the anterior hip joint can include:

- Abnormal hip or acetabular morphology (including FAI and dysplasia)
- Repetitive twisting sports such as tennis or golf
- Excessive anterior translation of the femoral head

To improve patient recovery following hip arthroscopy it is important to assess and address any pre-disposing factors that may have contributed to labral tear. This includes thorough assessment of the pelvis, hip and lower limb chain. Poor motor control strategies around the lumbar-pelvic and pelvic-hip region, and postural malalignment can all contribute to functional hip impingement or excessive anterior translation of the femoral head. When a labral tear presents as a traumatic injury there may be no pre-disposing factors evident.



Aims of Physiotherapy

- Minimise post-operative pain and swelling
- Restore hip and pelvic functional stability and strength
- Restore hip ROM
- Restore proprioception for return to work and sport
- Address pre-disposing factors contributing to biomechanical overload on anterior hip structures

Rehabilitation Program:

The rehabilitation program is not a recipe. Modifications and considerations should be made according to;

- Patient goals
- Patient age and pre-operative level of fitness
- Pre-existing contributing factors to labral tear
- Any associated concurrent dysfunction in the hip and pelvic region

Assessment and rehabilitation of the hip and pelvic area is a complex and requires thorough analysis of the pelvis, hip and lower limb. All times and exercises provided in this protocol are to serve as guidelines. Actual progress may be faster or slower depending on the individual patient.

An early goal is to restore a normal gait pattern based on symptoms and functional pelvic-hip control. For this reason it may be beneficial to use crutches in the first week only if a limp is present, until normal gait pattern is restored. For patients undergoing labral resection crutches are not usually required. For patients undergoing labral repair it is recommended a period of 2-3 weeks PWB on crutches.

Typically 4 to 6 weeks is needed to recover from the aspects of surgical intervention including intra-articular effusion. Therefore pushing end range of movement during this time can be counterproductive to a patient's recovery. Hip ROM is encouraged only to tolerance.

Throughout rehabilitation manual therapy by the physiotherapist (such as massage) may be appropriate for patients with a tendency to hypertonicity in deep posterior hip muscles, TFL, adductors or hamstrings. Manual therapy for lumbar or SIJ articular restriction may also be appropriate providing care is taken to avoid hip traction, end range hip flexion and end range internal rotation.

Gradual resumption of activities over a 6 to 16 week period is expected. During this time low-impact closed chain exercises can be progressed to a functional hip and pelvis rehabilitation program. Modifications to exercises should be considered according to the patients pre-disposing factors and addressing these factors is essential during this time frame. This will incorporate retraining motor patterns around the hip and pelvic region, which has more emphasis on motor control than power or strength of individual muscles. Rehabilitative exercises should not be painful within the hip joint or labrum. Return to sport will depend on the patient's pre-operative level of function and their ability to control the hip and pelvis in dynamic single leg balance. Stationary bike or cycling (if pain free) may be appropriate in the early stages of rehabilitation (2 to 6 weeks) if the patient wants to maintain cardiovascular fitness. Light jogging may be appropriate following the 8 to 12 week mark. Returning to hip challenges such as change of direction or speed work should be assessed on an individual basis.



Phase 1: 0-14 days

Goals:

- Reduce swelling and pain
- Restore normal gait pattern
- Restore normal standing and sitting postural alignment
- Regain ROM within tolerance

Treatment guidelines:

- Ice or NSAIDS as required
 - Crutches PWB 2-3 weeks for labral repair, and only if required for labral resection.
- Training on correct gait pattern
- Analysis and patient education on standing postural alignment to offload anterior hip. Improve patient awareness of correct standing hip and pelvic alignment (without posterior or lateral tilt)
 - Exercises to improve postural endurance, which should begin with inner unit and basic core contractions without substitution of more global muscles.
 - Start supine active assisted and active hip ROM exercises within tolerance. This should include patient awareness of maintaining lumbar-pelvic control and hip centred with accurate activation of lumbar-pelvic stability muscles (transversus, multifidus and pelvic floor). Care is to be taken with flexion and rotation ROM.
 - Start low load tasks such as quarter and half squats maintaining hip centering and alignment, core activation and body weight toward heels



Poor standing alignment with hip anterior of base of support



Hip ROM with core control



Squats with hip centered & core active



Phase 2: 2-6 weeks

Goals:

- Address any pre-disposing factors contributing to excessive load on anterior hip structures; including lumbar-pelvic stability, SIJ dysfunction, lower limb biomechanics.
- Restore hip- pelvic stability & strength with low load closed chain functional exercises.

Treatment guidelines:

- Assessment of lumbar-pelvic region. This can include:
Assessment and manual therapy treatment as appropriate for articular restriction in the lumbar spine or pelvis. Analysis of motor control around the lumbar spine and pelvis including accurate activation and timing of pelvic floor, multifidus and transversus, without substitution of more global muscles (such as external oblique or posterior pelvic floor)
- Assessment of the pelvic-hip region. This can include:
Analysis of dynamic hip stability including single leg balance (watching for excessive lateral shift, pelvic tilt, toe clawing). Analysis of hip motor recruitment patterns watching for signs of excessive use of global muscles including TFL and hamstrings.

Therapeutic exercise prescription in this phase will be determined by findings of lumbar-pelvic and hip assessment. As a general rule progression of exercises should always include activation of deep stability muscles and cues to allow centering of the femoral head. Once accurate activation can be maintained exercises can be progressed to include retraining of good functional movement patterns. Foot and knee alignment during these exercises must be also maintained. A focus on low load and higher repetitions will help retrain motor control pattern and endurance. Exercises can include:


- Supine core activation with hip movements (such as pilates matwork). Accurate activation of inner unit must be maintained along with hip centering. Lower limb support with the use of band (pictured) or ball can assist in reducing load through anterior hip
- Lunges with pelvic and hip control, maintaining foot and knee alignment and loading through heel to encourage centered alignment of femoral head
- Weight shift and transfer with pelvic and hip control in half squat, quarter squat and upright postures
- Step ups and step downs with pelvic, hip and knee control
- Addition of theraband around the knees to encourage gluts activation during squats, lunges and step ups



Lunges with hip control
Operated hip forward



Use of resistance band to supine hip
control exercises to support hip



Use of resistance band

Phase 3: 6-12 weeks

Goals:

- Restore proprioception
- Progress functional exercises to include more challenging pelvic and hip control work
- Continue addressing pre-disposing factors

Treatment guidelines

- Progress squats and lunges to include proprioceptive challenges such as air cushion or spin discs. Can include hand held weights to increase strengthening.
- Increase weight transfer exercises to balance work, whilst maintaining activation of inner unit muscles and pelvic-hip control
- Gradually increase gluteal challenges with the use of theraband (such as multi-directional leg movements in single leg balance). This is provided the patient can maintain accurate hip and pelvic alignment and is not substituting with over active TFL or hamstrings

Phase 4: 12 weeks plus

Goals:

- Improved functional strength and endurance
- Return to sporting activities

Treatment guidelines:

- Increase challenges to proprioception including single leg balance on air cushion or wobble board
- Sports specific retraining with hip and pelvic control



Squats and lunges with air cushion

Balance with theraband Balance with air cushion



Hip Arthroscopy: Guidelines for manual therapy and exercise

	Phase One Week 1 & 2	Phase Two Week 3- 6	Phase Three Week 6-12	Phase Four Week 12 +
Gait retraining	•	•		
Standing posture/sitting posture	•	•		
Hip ROM to tolerance	•	•		
Manual therapy				
Massage to:				
Piriformis	•	•	•	
TFL and ITB		•	•	
Adductors		•	•	
Joint mobilisation to lumbar if required		•	•	
Joint mobilisation to SIJ if required		•	•	
Stretches if required:				
Piriformis		•	•	
Gastrocs	•	•	•	
Quads		•	•	
Hamstrings		•	•	
Adductor			•	
Core exercises with hip centered:				
Supine pilates matwork	•	•	•	•
Supine reformer level one	•	•	•	•
Functional gluteal exercises:				
Performed with accurate core activation				
Squats	•	•	•	
Lunges		•	•	
Squats with theraband		•	•	•
Lunges with theraband		•	•	•
Weight transfer		•	•	•
Step ups		•	•	•
Step Downs			•	•
Lunges with hand weights			•	•
Proprioception				
Squats on air cushion			•	•
Lunges on air cushion			•	•
Balance			•	•
Balance with theraband challenges				•

